

| | Clearwater County | Idaho County | Latah County | Lewis County | Nez Perce County | State of Idaho | Top Us Performers Benchmark |
|---|-------------------|--------------|--------------|--------------|------------------|----------------|-----------------------------|
| HEALTH OUTCOMES | 39 | 23 | 8 | 3 | 33 | | |
| Mortality Length of Life | 41 | 31 | 16 | 10 | 37 | | |
| Premature death — Years of potential life lost before age 75 per 100,000 population (age-adjusted) | 10,163 | 7,414 | 5,943 | 5,240 | 8,594 | 6,046 | 5,200 |
| Morbidity Quality of Life | 27 | 14 | 5 | 2 | 17 | | |
| Poor or fair health — Percent of adults reporting fair or poor health (age-adjusted) | 20% | 21% | 12% | 14% | 15% | 15% | 10% |
| Poor physical health days — Average number of physically unhealthy days reported in past 30 days (age-adjusted) | 4.1 | 4 | 3.6 | 3.3 | 4 | 3.6 | 2.5 |
| Poor mental health days — Average number of mentally unhealthy days reported in past 30 days (age-adjusted) | 3.9 | 3 | 2.8 | 2.8 | 3.5 | 3.4 | 2.3 |
| Low birthweight — Percent of live births with low birthweight (< 2500 grams) | 5.4% | 5.1% | 5.1% | 3.8% | 5.8% | 6.5% | 5.9% |
| HEALTH FACTORS | 39 | 32 | 5 | 25 | 15 | | |
| Health Behaviors | 39 | 28 | 10 | 30 | 37 | | |
| Adult smoking — Percent of adults that report smoking >= 100 cigarettes and currently smoking | 23% | 18% | 14% | 16% | 20% | 17% | 14% |
| Adult obesity — Percent of adults that report a BMI >= 30 | 34% | 27% | 27% | 30% | 32% | 27% | 25% |
| Food Environment Index - measure ranging from 0 (worst) to 10 (best) which equally weights two indicators of the food env. (access) | 6.5 | 5.9 | 6.7 | 5.9 | 7.5 | 7.2 | 8.4 |
| Physical inactivity — Percent of adults age 20 and over reporting no leisure activity | 23% | 25% | 16% | 22% | 23% | 19% | 20% |
| Access to exercise opportunities — Percent of the population with adequate access to locations for physical activity | 47% | 59% | 79% | 50% | 79% | 79% | 92% |
| Excessive drinking — Binge plus heavy drinking | 14% | 19% | 22% | 28% | 19% | 15% | 10% |
| Alcohol-impaired driving deaths - percentage of motor vehicle crash deaths with alcohol involvement | 30% | 44% | 32% | 20% | 26% | 32% | 14% |
| Sexually transmitted infections — Chlamydia rate per 100,000 population | 116 | 86 | 272 | 309 | 402 | 285 | 138 |
| Teen birth rate — Teen birth rate per 1,000 female population, ages 15-19 | 40 | 34 | 10 | 29 | 32 | 35 | 20 |
| Clinical Care | 24 | 35 | 7 | 37 | 5 | | |
| Uninsured adults — Percent of population under age 65 without health insurance | 19% | 21% | 15% | 21% | 17% | 18% | 11% |
| Primary care physicians — Ratio of population to primary care physicians | 954:1 | 1631:1 | 1469:1 | 3889:1 | 1275:1 | 1618:1 | 1045:1 |
| Dentists — Ratio of population to dentists | 2144:1 | 2686:1 | 2240:1 | 1951:1 | 1478:1 | 1565:1 | 1377:1 |
| Mental Health providers - Ratio of population to mental health providers | 953:1 | 948:1 | 680:1 | 976:1 | 493:1 | 554:1 | 386:1 |
| Preventable hospital stays — Hospitalization rate for ambulatory-care sensitive conditions per 1,000 Medicare enrollees | 61 | 60 | 42 | 56 | 41 | 36 | 41 |
| Diabetic monitoring — Percent of diabetic Medicare enrollees that receive HbA1c screening | 80% | 79% | 87% | 73% | 86% | 82% | 90% |
| Mammography screening — Percent of female Medicare enrollees that receive mammography screening | 56.1% | 54.9% | 61.9% | 52.7% | 66.6% | 58.9% | 70.7% |
| Social & Economic Factors | 36 | 32 | 3 | 17 | 10 | | |
| High school graduation — Percent of ninth grade cohort that graduates in 4 years | 95% | 84% | 82% | 91% | 83% | 81% | |
| Some college — Percent of adults aged 25-44 years with some post-secondary education | 52.2% | 55.9% | 80.5% | 59.0% | 6670.0% | 63.7% | 71.0% |
| Unemployment — Percent of population age 16+ unemployed but seeking work | 12.1% | 8.8% | 5.7% | 5.3% | 5.4% | 6.2% | 4.0% |
| Children in poverty — Percent of children under age 18 in poverty | 26% | 23% | 15% | 25% | 18% | 19% | 13% |
| Income inequality - Ratio of household income at the 80th percentile to income at the 20th percentile | 3.5 | 3.7 | 4.9 | 4.0 | 3.8 | 4.1 | 3.7 |
| Children in single-parent households — Percent of children that live in household headed by single parent | 21% | 31% | 19% | 30% | 32% | 25% | 20% |
| Social associations - Number of membership associations per 10,000 population | 11.6 | 5.5 | 10.2 | 12.9 | 12.9 | 7.7 | 22.0 |
| Violent Crime Rate — Violent crime rate per 100,000 population | 201 | 108 | 120 | 139 | 151 | 210 | 59 |
| Injury deaths - Injury mortality per 100,000 | 87 | 107 | 56 | 94 | 100 | 66 | 50 |
| Physical Environment | 31 | 17 | 30 | 26 | 23 | | |
| Air pollution-particulate matter days — Average daily measure of fine particulate matter in micrograms per cubic meter | 10.9 | 10.4 | 11.1 | 10.7 | 10.7 | 10.1 | 9.5 |
| Drinking water safety —percentage of population exposed to water exceeding a violation limit during the past year | 5% | 11% | 2% | 8% | 2% | 6% | 0% |
| Severe housing problems - Percentage of households with at least 1 of 4 housing problems: overcrowding, high housing costs, or lack of kitchen or plumbing facilities | 10% | 14% | 20% | 13% | 12% | 16% | 9% |
| Driving alone to work - Percent of the workforce that drives alone to work | 81% | 72% | 67% | 75% | 81% | 78% | 71% |
| Long commute - driving alone - Among workers who commute in their car alone, the percentage that commute more than 30 minutes | 26% | 21% | 21% | 31% | 11% | 21% | 15% |

Note: Blank values reflect unreliable or missing data

New Measures